




Test de Bronchoprovocation au mannitol: médicaments à cesser avant le test

Mannitol Challenge test: Medications to withhold prior to testing

Minimum time to withhold	Medication
8 heures/ 8 hours	<p>Bêta-2 agonistes à courte durée d'action/ Short-acting beta-2 agonists: salbutamol (Ventolin®), Apo-salvent, Teva-salbutamol, terbutaline (Bricanyl®)</p> <div style="display: flex; justify-content: space-around; align-items: center;">      </div> <p>Anti-inflammatoires non stéroïdiens inhalés/ Inhaled non-steroidal anti-inflammatory cromolyn sodium (Intal®), nedocromil sodium (Tilade®)</p> <div style="display: flex; justify-content: space-around; align-items: center;">   </div>

<p>24 heures/ 24 hours</p>	<p>Corticostéroïdes inhalés / Inhaled corticosteroids budesonide (Pulmicort®), ciclesonide (Alvesco®) fluticasone (Flovent®), mometasone (Asmanex®), beclomethasone (Qvar®)</p>   <p>Anticholinergiques à action prolongée / Anticholinergic Bronchodilators: ipratropium bromide (Atrovent®)</p> 
<p>36 heures/ 36 hours</p>	<p>Bêta- 2 agonistes de longue durée d'action / Long –acting Beta2- agonists salmeterol (Serevent®), formoterol (Oxeze®)</p>  <p>Combinaison de corticostéroïdes inhalés et de bêta-2- agonistes à action prolongée/ Combined Inhaled corticosteroids and long –acting Beta2 agonists</p> <p>fluticasone and salmeterol (Advair®), budesonide and formoterol (Symbicort®), mometasone and formoterol (Zenhale®)</p> 

48 heures/ 48 hours	<p>Combinaison de corticostéroïdes inhalés et de bêta-2-agonistes à ultra longue durée d'action /</p> <p>Combined inhaled corticosteroid and and ultra-long-acting beta-2 agonist:</p> <p>fluticasone and vilanterol (Breo Ellipta®)</p> 
72 heures/ 72 hours Pilules / Pills	<p>Anticholinergiques à action prolongée / Long –Acting Anticholinergics:</p> <p>tiotropium bromide (Spiriva®)</p>  <p>Antihistaminiques/ Antihistamines:</p> <p>loratadine (Claritin®), certirizine (Zyrtec®) (Reactine®), fexofenadine (Allegra®), brompheniramine (Dimetapp®), levocetirizine (Xyzal®), desloratadine (Aerius®), diphenhydramine (Benadryl®)</p>
4jours/ 4 days Pilules / Pills	<p>Antagonistes des récepteurs des leucotriènes/ Leukotriene-Receptor Antagonists</p> <p>montelukast (Singulair®), zafirlukast (Accolate®)</p>  