

## CRITERIA FOR NEONATAL FOLLOW-UP

The Neonatal Follow-up Clinic follows patients discharged from the NICU, who have risk factors for neurodevelopmental delay; the following are **guidelines** for referral.

### CORE CRITERIA

These criteria are directly related to the expertise of our level III NICU.

- Prematurity: gestational age <29 weeks
- Asphyxia/Hypoxic-Ischemic Encephalopathy (HIE)
  - Moderate (Modified Sarnat 2) and severe (Modified Sarnat 3), with or without cooling (use worst recorded score)
  - Mild (Modified Sarnat 1): if abnormal neurodevelopmental exam or imaging or if in a study (modified follow-up protocol)
- Broncho-Pulmonary Dysplasia (BPD) oxygen dependent
- Post Extracorporeal Membrane Oxygenation (ECMO)
- Status post early open heart surgery (< 3 months of age)
- Home Enteral Feeding Program (HEFP) at discharge from NICU, not related to a gastrointestinal pathology followed by GI or by the Complex care team

### LOCAL CRITERIA

- MUHC “Inborn” triplets

### PROGRAM VISIT SEQUENCE: baseline scheduled protocol

#### ➤ Core Criteria

By age (corrected age, if applicable):

- 4 m, 9 m, 18 m, 36 m and preschool (in year prior to kindergarten entry age 5)
- Subsequent visit and extra visits possible, on clinical basis
- For HEFP: Follow up could be discontinued, if patient off gavage with no other criteria

### OTHER CRITERIA

#### **ANTEPARTUM AND DELIVERY**

- Intrauterine growth restriction (IUGR): Birth weight < 3 standard deviations (see tables)

#### **NEUROLOGIC**

- Neonatal seizures
- Microcephaly (birth head circumference < 3%)

- Intraventricular hemorrhage grade 3 and 4
- Sensory deficits (visual, auditory), including newborns referred by the Universal Hearing Screening Program
- Abnormal neurodevelopmental exam (at discharge, if hospitalization > 48 hrs)
- Abnormal Significant radiologic findings: *Parenchymal cerebral lesions, persistent ventriculomegaly, hypoxic-ischemic changes, periventricular leukomalacia, significant subarachnoid or subdural hemorrhage*

## **INFECTION**

- Meningitis (with or without positive cultures)
- Congenital infections (TORCH)

## **ADDITIONAL**

- Multiple congenital anomalies, undiagnosed syndrome
- Certain genetic syndromes associated with neurodevelopmental delay (e.g. Trisomy 13, 18, Rubenstein Taybi, DiGeorge, CHARGE)
- Severe hemodynamic compromise (hypovolemic/septic shock)
- Study (modified follow-up as required)
- Other exceptional cases, as discussed with the NNFU team

## **PROGRAM VISIT SEQUENCE**

### ➤ **Other Criteria**

#### By age:

- 4 m, 9 m, 18 m and 36 m

#### After 36 mos:

- If development normal, discharge. File could be re-opened on clinical basis
- If abnormal neurodevelopmental assessment, consider to pursue Follow-up visit
- If child is followed in a rehabilitation center, consider discharge from Neonatal program

We accept referrals of children who meet our “CORE” criteria and are under 5 years of age (children not born at the MUHC).

- A consultation form is needed/with summary, if possible

Therapists from our multidisciplinary team (OT/PT/Speech/Audio/Nutrition/Psychology/Social Work) may also refer children under 18 months of age (<18 mo) for neurodevelopmental concerns, questions/needs. A consult request is needed.